arXiv:1611.04130v1 [math.QA] 13 Nov 2016

NONCOMMUTATIVE BORSUK-ULAM-TYPE CONJECTURES REVISITED

LUDWIK DĄBROWSKI, PIOTR M. HAJAC, AND SERGEY NESHVEYEV

Abstract. Let H be the C*-algebra of a non-trivial compact quantum group acting freely on a unital C*-algebra A. Baum, Dąbrowski and Hajac conjectured that there does not exist an equivariant $*$-homomorphism from A to the equivariant noncommutative join C*-algebra $A \circledast^{\delta} H$. When A is the C*-algebra of functions on a sphere, and H is the C*-algebra of functions on $\mathbb{Z}/2\mathbb{Z}$ acting antipodally on the sphere, then the conjecture becomes the celebrated Borsuk-Ulam theorem. Recently, Chirvasitu and Passer proved the conjecture when H is commutative. In a simple way, we extend this result to a far more general setting assuming only that H admits a character different from the counit. We show how our result implies a noncommutative Brouwer fixed-point theorem and, in particular, the non- contractibility of such compact quantum groups. Moreover, assuming that our compact quantum group is the q-deformation of a compact connected semisimple Lie group, we prove that there exists a finite-dimensional representation of the compact quantum group such that, for any C*-algebra A admitting a character, the finitely generated projective module associated with $A \circledast^{\delta} H$ via this representation is not stably free.

Contents

1. Introduction 2
1.1. Around the classical Borsuk-Ulam 2
1.2. Quantum preliminaries 3
2. The noncommutative Borsuk-Ulam conjecture of type I 5
2.1. The pointed noncommutative Borsuk-Ulam theorem of type I 5
2.2. Noncommutative Brouwer fixed-point theorem and non-contractibility 6
3. The noncommutative Borsuk-Ulam conjecture of type II 6
3.1. The pointed noncommutative Borsuk-Ulam theorem of type II 6
3.2. The K-theoretic noncommutative Borsuk-Ulam theorem for the q-deformations of some compact Lie groups 7
Acknowledgements 8
References 8

1. Introduction

The goal of this paper is to prove, under some additional assumptions, both types of the conjecture stated in [2, Conjecture 2.3]. We will also derive some consequences of the type I conjecture, and a K-theoretic strengthening of the type II conjecture.

1.1. **Around the classical Borsuk-Ulam.** The Borsuk-Ulam theorem [4] is a fundamental theorem of topology concerning spheres in Euclidean spaces. It can be phrased in the following three equivalent ways:

Theorem 1.1 (Borsuk-Ulam). *For any natural number n, if $f\colon S^{n+1}\to\mathbb{R}^{n+1}$ is continuous, then there exists a pair $(p,-p)$ of antipodal points on S^{n+1} such that $f(p)=f(-p)$.*

Theorem 1.2 (equivariant formulation). *For any natural number n, there does* not *exist a $\mathbb{Z}/2\mathbb{Z}$-equivariant continuous map $\widetilde{f}\colon S^{n+1}\to S^n$.*

Theorem 1.3 (join formulation). *For any natural number n, there does* not *exist a $\mathbb{Z}/2\mathbb{Z}$-equivariant continuous map $\widetilde{f}\colon S^n*\mathbb{Z}/2\mathbb{Z}\to S^n$.*

The third formulation of the theorem lends itself to a natural and wide generalization, which was proposed in [2, Conjecture 2.2], and recently proved in [5].

Theorem 1.4 ([5]). *Let X be a compact Hausdorff space equipped with a continuous free action of a non-trivial compact Hausdorff group G. Then, for the diagonal action of G on the join $X*G$, there does* not *exist a G-equivariant continuous map $X*G\to X$.*

By taking any $x_0\in X$, we can define a G-equivariant continuous map $G\ni g\mapsto x_0g\in X$. Consequently, we obtain:

Corollary 1.5. *There does* not *exist a G-equivariant continuous map $X*G\to G$. In particular, for $X=G$ this means that G is not contractible.*

In the case G is a compact connected semisimple Lie group, the above corollary also follows from:

Theorem 1.6. *Let G be a compact connected semisimple Lie group. Then there exists a finite-dimensional representation V of G such that for any compact Hausdorff space X equipped with a free G-action, the associated vector bundle*

$$(X*G)\overset{G}{\times}V$$

is not *stably trivial.*

Proof. To begin with, let us show that it suffices to prove the theorem in the special case $X=G$. Indeed, let Y and Z be compact Hausdorff G-spaces. Assume that the G-action on Z is free, then the G-action on Y is automatically free as well. If $f:Y\to Z$ is a continuous G-equivariant map, then it induces a continuous map between quotient spaces $\bar f:Y/G\to Z/G$. Now, for any finite-dimensional representation $G\to\mathrm{GL}(V)$, we obtain:

$$\bar f^*\colon K^0(Z/G)\ni\left[Z\overset{G}{\times}V\right]\longmapsto\left[Y\overset{G}{\times}V\right]\in K^0(Y/G). \tag{1.1}$$

Hence, the stable non-triviality of the associated vector bundle $Y \times^G V$ implies the stable non-triviality of the associated vector bundle $Z \times^G V$.

Finally, since for any $x_0 \in X$ the continuous G-equivariant map $G \ni g \mapsto x_0 g \in X$ induces a continuous G-equivariant map $G * G \to X * G$, the forgoing argument proves that it suffices to prove the theorem for $X = G$. Therefore, the following lemma concludes the proof. □

LEMMA 1.7. *Let G be a compact connected semisimple Lie group. Then there exists a representation of G on a finite-dimensional vector space V such that*

$$[(G * G) \times^G V] \neq [\dim V] \quad \textit{in} \quad K^0(\mathcal{S}G).$$

Here $\mathcal{S}G$ is the unreduced suspension of G, which we identify with $(G * G)/G$.

Proof. Put $E_V := (G * G) \times^G V$ for brevity. Consider the group

$$\tilde{K}^0(\mathcal{S}G) = \operatorname{Ker}\big(K^0(\mathcal{S}G) \longrightarrow K^0(\mathrm{pt}) = \mathbb{Z}\big), \tag{1.2}$$

where the homomorphism is defined by any point of the connected space $\mathcal{S}G$. As was already observed in [1], using the description of $K^1(G)$ in terms of homotopy classes of maps $G \to \mathrm{GL}(\infty)$, it is not difficult to see that there is an isomorphism $\beta\colon \tilde{K}^0(\mathcal{S}G) \to K^1(G)$ such that $\beta([\dim V] - [E_V])$ is exactly $[u] \in K^1(G)$, where $u : G \to \mathrm{GL}(V)$ is the representation defining E_V. Indeed, by specifying [2, Theorem 3.3] to the classical situation, and identifying χ^{-1} in that theorem with u, we see that $\beta^{-1}([u]) = [\dim V] - [E_V]$, where β^{-1} is the C*-algebraic (topological) version of the Milnor connecting homomorphism [10, Section 0.4].

Now, if G is simply connected, then for any fundamental representation V we have $\beta([\dim V] - [E_V]) \neq 0$ by [11]. If G is not simply connected, it is a quotient of its simply connected cover $\widetilde{G}$ by a finite central subgroup Γ. Therefore, to get a required representation of the quotient $G = \widetilde{G}/\Gamma$, it suffices to consider the tensor power $V^{\otimes d}$ for a fundamental representation V of $\widetilde{G}$, where $d = |\Gamma|$. Then, as $K^1(\widetilde{G})$ has no torsion [11], the $K^1(\widetilde{G})$-class

$$\beta\big(\big[\dim V^{\otimes d}\big] - \big[E_{V^{\otimes d}}\big]\big) = d(\dim V)^{d-1}\beta\big(\big[\dim V\big] - \big[E_V\big]\big)$$

(see [11, Section I.4]) is nonzero, whence $\beta\big(\big[\dim V^{\otimes d}\big] - \big[E_{V^{\otimes d}}\big]\big)$ viewed as a class in $K^1(G)$ is also nonzero. □

1.2. **Quantum preliminaries.** In this section, we review basic noncommutative concepts used to formulate quantum Borsuk-Ulam-type statements. We always assume that *-homomorphisms between unital C*-algebras are unital.

1.2.1. *Compact quantum groups* [16]. Recall that a compact quantum group (H, Δ) consists of a unital C*-algebra H and a *coproduct* $\Delta : H \to H \otimes_{\min} H$ that is an injective coassociative $*$-homomorphism enjoying *cancellation laws*:

$$\{\Delta(g)(1 \otimes h) \,|\, g, h \in H\}^{cls} = H \otimes_{\min} H = \{(g \otimes 1)\Delta(h) \,|\, g, h \in H\}^{cls}. \tag{1.3}$$

Here *cls* stands for "closed linear span".

1.2.2. *Free actions* [15, 8]. Let A be a unital C*-algebra, and (H,Δ) be a compact quantum group. We say that the quantum group (H,Δ) *acts* on A, iff there is an injective $*$-homomorphism $\delta : A \to A\otimes_{\min} H$ (called a *coaction*) satisfying:

(1) $(\delta\otimes\mathrm{id})\circ\delta = (\mathrm{id}\otimes\Delta)\circ\delta$ (coassociativity),
(2) $\{\delta(a)(1\otimes h)\,|\,a\in A, h\in H\}^{cls} = A\otimes_{\min} H$ (counitality).

We call such an action *free*, iff $\{(x\otimes 1)\delta(y)\,|\,x,y\in A\}^{cls} = A\otimes_{\min} H$.

1.2.3. *Equivariant join* [7, 3]. There is an obvious definition of the join of unital C*-algebras

$$A\circledast B := \left\{ f\in C\big([0,1], A\underset{\min}{\otimes} B\big) \;\middle|\; f(0)\in\mathbb{C}\otimes B,\ f(1)\in A\otimes\mathbb{C}\right\} \tag{1.4}$$

dualizing the join of compact Hausdorff spaces: $C(X)\circledast C(Y) = C(X*Y)$. In particular, we can take $B=H$, where (H,Δ) is a compact quantum group acting on A.

However, to make this construction equivariant as in the classical case, we need to modify the above definition to become:

Definition 1.8. *Let (H,Δ) be a compact quantum group acting on a unital C*-algebra A via $\delta : A\to A\otimes_{\min} H$. We call the unital C*-algebra*

$$A\overset{\delta}{\circledast} H := \left\{ f\in C\big([0,1], A\underset{\min}{\otimes} H\big) \;\middle|\; f(0)\in\mathbb{C}\otimes H,\ f(1)\in\delta(A)\right\} \tag{1.5}$$

the equivariant noncommutative join *of A and H.*

The coaction

$$(\mathrm{id}\otimes\Delta)\circ : C\big([0,1], A\underset{\min}{\otimes} H\big) \longrightarrow C\big([0,1], A\underset{\min}{\otimes} H\big)\underset{\min}{\otimes} H$$

induces a free action of (H,Δ) on $A\circledast^{\delta} H$.

1.2.4. *The Peter-Weyl subalgebra and associated modules* [3]. Let A be a unital C*-algebra with a free action of a compact quantum group (H,Δ) given by a coaction $\delta : A\to A\otimes_{\min} H$. We define the *Peter-Weyl subalgebra* of A (see [3]) as

$$\mathcal{P}_H(A) := \{\, a\in A \mid \delta(a)\in A\otimes\mathcal{O}(H)\,\}. \tag{1.6}$$

Here $\mathcal{O}(H)$ is the Hopf algebra spanned by the matrix coefficients of irreducible unitary corepresentations of H. Using the coassociativity of δ, one can check that $\mathcal{P}_H(A)$ is a right $\mathcal{O}(H)$-comodule algebra. For $(A,\delta)=(H,\Delta)$ we have $\mathcal{P}_H(H)=\mathcal{O}(H)$.

Now we can define the cotensor product

$$\mathcal{P}_H(A)\Box V := \{t\in\mathcal{P}_H(A)\otimes V \mid (\delta\otimes\mathrm{id})(t) = (\mathrm{id}\otimes {}_V\Delta)(t)\}, \tag{1.7}$$

where ${}_V\Delta\colon V\to\mathcal{O}(H)\otimes V$ is the given left coaction of $\mathcal{O}(H)$ on V. We know from [3] that if $\dim V<\infty$, then the cotensor product $\mathcal{P}_H(A)\Box V$ is finitely generated projective as a left module over the fixed-point subalgebra $A^{\mathrm{co}H} := \{a\in A\mid\delta(a)=a\otimes 1\}$.

Next, we unravel the meaning of the thus defined associated module for commutative C*-algebras $A=C(X)$ and $H=C(G)$ (the classical setting). Let $\varrho\colon G\to\mathrm{GL}(V)$

be a representation of G on a finite-dimensional vector space V. We view V as a left $\mathcal{O}(G)$-comodule and denote by ${}_V\Delta\colon V \to \mathcal{O}(G)\otimes V$ the corresponding map, given by the formula $({}_V\Delta(v))(g) = \varrho(g^{-1})v$. note that the $C(X/G)$-module $C_G(X,V)$ of all continuous G-equivariant functions from X to V is naturally isomorphic with the $C(X/G)$-module $\Gamma(X\times^G V)$ of all continuous sections of $X\times^G V$. Here G-equivariance means

$$\forall\, x\in X,\, g\in G:\ f(xg) = \varrho(g^{-1})(f(x)). \tag{1.8}$$

Combining this with [3, Corollary 4.1], we can naturally identify $\mathcal{P}_{C(G)}(C(X))\square V$ with $\Gamma(X\times^G V)$ as left $C(X/G)$-modules.

2. The noncommutative Borsuk-Ulam conjecture of type I

Let us begin by recalling the noncommutative Borsuk-Ulam conjecture of type I.

Conjecture 2.1 ([2]). *Let A be a unital C*-algebra equipped with a free action of a non-trivial compact quantum group (H,Δ), and let $A\circledast^{\delta} H$ be the equivariant noncommutative join C*-algebra of A and H with the induced free action of (H,Δ). Then*

$$\nexists \text{ an } H\text{-equivariant } *\text{-homomorphism } A\to A\circledast^{\delta} H. \tag{2.9}$$

2.1. The pointed noncommutative Borsuk-Ulam theorem of type I.

Theorem 2.2. *Let A be a unital C*-algebra with a free action of a non-trivial compact quantum group (H,Δ), and let $A\circledast^{\delta} H$ be the equivariant noncommutative join C*-algebra of A and H with the induced free action of (H,Δ). Then, if H admits a character that is not convolution idempotent, the following equivalent statements hold:*

$$\nexists \text{ an } H\text{-equivariant } *\text{-homomorphism } A\longrightarrow A\circledast^{\delta} H, \tag{2.10}$$

$$\nexists\ *\text{-homomorphism } \gamma : A\longrightarrow \mathcal{C}A \text{ such that } \mathrm{ev}_1\circ\gamma : A\to A \text{ is } H\text{-colinear.} \tag{2.11}$$

Here $\mathcal{C}A := A\circledast\mathbb{C}$ is the cone of A, and ev_1 is the evaluation at 1.

Proof. Recall that a compact quantum group (K,Δ) is a closed quantum subgroup of (H,Δ), iff there is a surjective $*$-homomorphism $\pi\colon H\to K$ respecting the coproducts. In particular, when K is commutative, the induced coaction $(\mathrm{id}\otimes\pi)\circ\delta : A\to A\otimes_{\min} K$ defines an action of (K,Δ) on A, and consequently on $A\circledast^{(\mathrm{id}\otimes\pi)\circ\delta} K$. Furthermore, if $\varphi\colon A\to A\circledast^{\delta} H$ is an H-equivariant $*$-homomorphism, then $(\mathrm{id}\otimes\pi)\circ\varphi$ is a K-equivariant $*$-homomorphism $A\to A\circledast^{(\mathrm{id}\otimes\pi)\circ\delta} K$. Therefore, it suffices to prove (2.10) for (K,Δ) with K commutative.

Since (2.10) is proven in [5, Theorem 3.2] for any commutative $K\neq\mathbb{C}$, all we need to do is to show that we have such a commutative K. It follows from our assumption that H admits a character that is not convolution idempotent that the compact Hausdorff group G_H of characters of the C*-algebra H, equipped with the weak * topology and convolution product, is not a trivial group. Now the proof of (2.10) is concluded by taking $K := C(G_H)$ in [5, Theorem 3.2].

Finally, if $\gamma : A\to\mathcal{C}A$ is as in (2.11), then $\varphi := (\gamma\otimes\mathrm{id})\circ\delta$ descends to a map as in (2.10). Vice versa, if φ is as in (2.10), and if ε_H is the counit on H corresponding to

the unit of the classical subgroup G_H (note that here we do not need the non-triviality of G_H), then $\gamma := (\mathrm{id}\otimes\mathrm{id}\otimes\varepsilon_H)\circ\varphi$, descends to a map as in (2.11). □

Conjecture 2.1 remains open for compact quantum groups with simple C*-algebras, e.g. the reduced C*-algebra $C^*_{red}F_2$ of the free group on two elements.

2.2. **Noncommutative Brouwer fixed-point theorem and non-contractibility.**

2.2.1. *Noncommutative Brouwer fixed-point theorem.* As a corollary of (2.11), we obtain a noncommutative generalization of the Brouwer fixed-point theorem:

$$\nexists \text{ unital } *\text{-homomorphism } \gamma : A \longrightarrow \mathcal{C}A \text{ such that } \mathrm{ev}_1\circ\gamma = \mathrm{id}_A. \tag{2.12}$$

Indeed, in the particular case when $A = C(S^n)$ and $H = C(\mathbb{Z}/2\mathbb{Z})$, this corollary is well known to be equivalent to the Brouwer fixed-point theorem for continuous maps $B^n \to B^n$ between balls (see also [6]).

2.2.2. *Non-contractibility.* Theorem 2.2 restricted to the case when $(A,\delta) = (H,\Delta)$, boils down to the non-triviality of the principal H-bundle $H \circledast^{\Delta} H$. While in the classical setting $H = C(G)$ this is equivalent to the well-known [12] non-contractibility of non-trivial compact Hausdorff groups, in the quantum case the non-triviality of $H \circledast^{\Delta} H$ was known so far for the quantum group $SU_q(2)$ [2]. Now it follows for the q-deformations of any compact connected semisimple Lie group from Theorem 3.2, and for any compact quantum group whose C*-algebra admits a character that is not convolution idempotent from Theorem 2.10.

Finally, since the retraction of $\mathcal{C}A$ to A can be literally taken as a definition of *contractibility* of A, not only do we infer the non-contractibility of such compact quantum groups, but also we recover the classical mechanism relating the non-triviality of $G * G$ with the non-contractibility of G

3. The noncommutative Borsuk-Ulam conjecture of type II

While the noncommutative Borsuk-Ulam conjecture of type I remains open as stated, the noncommutative Borsuk-Ulam conjecture of type II requires corrections to avoid counterexamples. However, with additional assumptions, not only do we prove the original conjecture, but also its K-theoretic strengthening.

3.1. **The pointed noncommutative Borsuk-Ulam theorem of type II.**

The following result is an immediate corollary of Theorem 2.2. Under some additional assumptions, it proves the noncommutative Borsuk-Ulam conjecture of type II (see [2, Conjecture 2.3]).

Theorem 3.1. *Let A be a unital C*-algebra with a free action of a non-trivial compact quantum group (H,Δ), and let $A\circledast^{\delta} H$ be the equivariant noncommutative join C*-algebra of A and H with the induced free action of (H,Δ). Then, if H admits a character that is not convolution idempotent and A admits any character,*

$$\nexists \text{ an } H\text{-equivariant } *\text{-homomorphism } H \to A\circledast^{\delta} H. \tag{3.13}$$

Proof. If A has a character α, then an H-equivariant *-homomorphism as in (3.13) composed with the H-equivariant *-homomorphism $(\alpha\otimes\mathrm{id})\circ\delta$ would provide an H-equivariant *-homomorphism as in (2.10). Thus (2.10) implies (3.13). □

Assuming that A admits a character removes the counter-example [5, Theorem 3.6]) and affords a proof of (3.13), but we do not know when this assumption is necessary (see [5, Theorem 3.5]).

3.2. **The K-theoretic noncommutative Borsuk-Ulam theorem for the q-deformations of some compact Lie groups.** Let G be a compact connected (but not necessarily simply connected) semisimple Lie group. For $q > 0$ consider the quantized universal enveloping algebra $U_q\mathfrak{g}$, where $\mathfrak{g}$ is the Lie algebra of G. For every finite-dimensional representation ρ of G on V, there is a representation ρ_q of $U_q\mathfrak{g}$ on the same space determined uniquely, up to isomorphism, by the requirement that the dimensions of the weight spaces remain the same as for the original representation. The matrix coefficients of these representations span a Hopf *-subalgebra $\mathcal{O}(G_q)$ of $(U_q\mathfrak{g})^*$. The universal C*-completion of $\mathcal{O}(G_q)$ defines a compact quantum group $(C(G_q),\Delta)$. For every (V,ρ) as above, the representation ρ_q defines a left $\mathcal{O}(G_q)$-comodule structure on the vector space V. We denote the thus obtained comodule by V_q.

THEOREM 3.2. *Let G be a compact connected semisimple Lie group and V be as in Theorem 1.6. Then, for any $q > 0$ and any unital C*-algebra A admitting a character and equipped with a free action of $(C(G_q),\Delta_q)$, the associated finitely generated projective left $(A \circledast^{\delta} C(G_q))^{\mathrm{co}\,C(G_q)}$-module*

$$\mathcal{P}_{C(G_q)}\Big(A \overset{\delta}{\circledast} C(G_q)\Big)\Box V_q$$

is not *stably free.*

Proof. As in the classical case, we first reduce the theorem to the case $A = C(G_q)$. We use a character α on A to define a G_q-equivariant $*$-homomorphism $(\alpha\otimes\mathrm{id})\circ\delta : A \to C(G_q)$, which induces a G_q-equivariant $*$-homomorphism

$$f : A \overset{\delta}{\circledast} C(G_q) \longrightarrow C(G_q) \overset{\Delta}{\circledast} C(G_q).$$

Due to the equivariance, the $*$-homomorphism f restricts and corestricts to a $*$-homomorphism between fixed-point subalgebras:

$$f| : \Big(A \overset{\delta}{\circledast} C(G_q)\Big)^{\mathrm{co}\,C(G_q)} \longrightarrow \Big(C(G_q) \overset{\Delta}{\circledast} C(G_q)\Big)^{\mathrm{co}\,C(G_q)}.$$

Now we can apply [9, Theorem 2.1]) to conclude that we have

$$(f|)_* : K_0\Big(\big(A \overset{\delta}{\circledast} C(G_q)\big)^{\mathrm{co}\,C(G_q)}\Big) \longrightarrow K_0\Big(\big(C(G_q) \overset{\Delta}{\circledast} C(G_q)\big)^{\mathrm{co}\,C(G_q)}\Big),$$

$$(f|)_*\Big(\Big[\mathcal{P}_{C(G_q)}\big(A \overset{\delta}{\circledast} C(G_q)\big)\Box V_q\Big]\Big) = \Big[\mathcal{P}_{C(G_q)}\big(C(G_q) \overset{\Delta}{\circledast} C(G_q)\big)\Box V_q\Big].$$

Hence, if the module on the right-hand side is not stably free, also the module on the left-hand side is not stably free. Thus we have reduced the theorem to the special case $A = C(G_q)$, as desired.

Next, put

$$\Sigma C(G_q) := \Big(C(G_q) \overset{\Delta}{\circledast} C(G_q)\Big)^{\mathrm{co}\, C(G_q)}, \quad \Gamma(E_{V_q}) := \mathcal{P}_{C(G_q)}\Big(C(G_q) \overset{\Delta}{\circledast} C(G_q)\Big)\square V_q\,,$$

and observe that for $q = 1$ we have $\Sigma C(G) = C(\mathcal{S}G)$. Now, using the structure of a KK-fibration on the field of C*–algebras $C(G_q)$ defined in [13, 14], one can show that there exists an isomorphism

$$K_0(\Sigma C(G_q)) \ni \Gamma(E_{V_q}) \longmapsto \Gamma(E_V) \in K_0(C(\mathcal{S}G)).$$

Combining it with the Serre-Swan isomorphism

$$K_0(C(\mathcal{S}G)) \ni \Gamma(E_V) \longmapsto E_V \in K^0(\mathcal{S}G),$$

and using Lemma 1.7 we conclude the proof of the theorem. □

Acknowledgements

This work is part of the project *Quantum Dynamics* sponsored by EU-grant RISE 691246 and Polish Government grant 317281. This work was also partially supported by the European Research Council under the EU's Seventh Framework Programme (FP/2007-2013)/ERC Grant Agreement 307663. Finally, Ludwik Dąbrowski is grateful for his support at IMPAN provided by Simons-Fundation grant 346300 and a Polish Government MNiSW 2015–2019 matching fund.

References

[1] M.F. Atiyah: *On the K-theory of compact Lie groups*, Topology 4 (1965) 95–99.

[2] P.F. Baum, L. Dąbrowski, P.M. Hajac: *Noncommutative Borsuk-Ulam-type conjectures*, Banach Center Publications 106 (2015) 9–18; arXiv:1502.05756.

[3] P.F. Baum, K. De Commer, P.M. Hajac: *Free actions of compact quantum group on unital C*-algebras*, arXiv:1304.2812.

[4] K. Borsuk: *Drei Sätze über die n-dimensionale euklidische Sphäre*, Fund. Math. 35 (1933) 177–190.

[5] A. Chirvasitu, B. Passer: *Compact group actions on topological and noncommutative joins*, arXiv:1604.02173v2.

[6] L. Dąbrowski: *Towards a noncommutative Brouwer fixed-point theorem*, J. Geom. Phys. 105 (2016) 60–65.

[7] L. Dąbrowski, T. Hadfield, P.M. Hajac: *Equivariant join and fusion of noncommutative algebras*, SIGMA 11 (2015) 082.

[8] D.A. Ellwood: *A new characterisation of principal actions*, J. Funct. Anal. 173 no. 1 (2000) 49–60.

[9] P.M. Hajac, T. Maszczyk: *Pulling back noncommutative associated vector bundles and constructing quantum quaternionic projective spaces* arXiv:1601.00021.

[10] P.M. Hajac, A. Rennie, B. Zieliński: *The K-theory of Heegaard quantum lens spaces*, Journal of Noncommutative Geometry, 7 (2013) 1185–1216.

[11] L. Hodgkin: *On the K-theory of Lie groups*, Topology 6 (1967) 1–36.

[12] B. Hoffmann: *A compact contractible topological group is trivial*, Archiv Math. 32, no. 1 (1979) 585–587.

[13] S. Neshveyev, L. Tuset: *K-homology class of the Dirac operator on a compact quantum group*, Doc. Math. 16 (2011) 767–780.

[14] S. Neshveyev, L. Tuset: *Quantized algebras of functions on homogeneous spaces with Poisson stabilizers*, Comm. Math. Phys. 312 (2012) 223–250.

[15] P. Podleś: *Symmetries of quantum spaces. Subgroups and quotient spaces of quantum $SU(2)$ and $SO(3)$ groups*, Comm. Math. Phys. 170 (1995) 1–20.

[16] S.L. Woronowicz: *Compact quantum groups*, in *Symétries quantiques (Les Houches, 1995)*, 845–884, North-Holland, Amsterdam, 1998.

SISSA (Scuola Internazionale Superiore di Studi Avanzati), Via Bonomea 265, 34136 Trieste, Italy

E-mail address: dabrow@sissa.it

Instytut Matematyczny, Polska Akademia Nauk, ul. Śniadeckich 8, 00-656 Warszawa, Poland,

E-mail address: pmh@impan.pl

Department of Mathematics, University of Oslo, P.O. Box 105, 3 Blindern, 0316 Oslo, Norway,

E-mail address: sergeyn@math.uio.no